



TUBE WITH MEASURING STICK AND TUBE WINDER

1. Tap the tube upside down a few times to ensure all the cream is near the opening.
2. Attach the measuring stick to the tube.
3. Squeeze the tube to dispense the cream.
Tip: The plunger of the measuring stick will extend to show how much of the cream is dispensed. Continue squeezing until the correct dosage shows.
4. Detach the measuring stick from the tube and replace cap.
5. Press the plunger down to dispense the cream from the measuring stick.
6. Apply cream **only on the area of the skin directed by your prescriber.**
7. Clean the measuring stick with warm water after each use.
8. Attach tube winder as needed to completely use all cream in tube.



TOPI-CLICK™

1. Remove the cap by placing thumb under extended lip on the cap and pushing upward.
2. Tap on hard surface to settle cream before dispensing.
3. Turn the base counterclockwise to dispense 1/4 gram per click.
Tip: Repeat step 3 as needed if the dosage prescribed by your physician is more than the dosage dispensed in one click.
4. Apply cream by rubbing applicator pad **only on the area of skin specified by your prescriber.**
5. Replace cap.
Note: Topi-Clicks™ are not refillable.
6. Store upright.



TOPI-PUMP™ AND SLEEK TOPI-PUMP™

1. Remove the cap.
2. Tap on hard surface to settle cream before dispensing.
3. Press the top of the pump down completely to dispense 1/2 gram for the Topi-Pump™ or 1/4 gram for the Sleek Topi-Pump™.
Tip: Repeat step 3 as needed if prescribed dosage is more than the amount dispensed in one pump.
4. Apply cream **only on the area of the skin directed by your prescriber.**
5. Replace cap.
Note: Topi-Pumps™ and Sleek Topi-Pumps™ are not refillable.
6. Store upright.

HOW MUCH CREAM SHOULD YOU USE?



Please refer to this image according to the dosage your prescriber has indicated on your medication. If you have any questions, you can contact our customer service representatives at 888-222-2956.



ACUPEN™

1. Remove the cap.
2. Unlock the device by turning the endcap counter-clockwise.
3. Push the end of the AcuPen™ (the same way you push the end of an ink pen) to dispense 1/8 gram.
Tip: Repeat step 3 as needed if prescribed dosage is more than the amount dispensed in one push.
4. Apply cream **only on the area of the skin directed by your prescriber.**
5. Lock the AcuPen™ by turning the end cap clockwise.
6. Replace cap.
Note: AcuPens™ are not refillable.



TRITURATE

- Small round compressed tablet.
- Custom options available upon request.
- Dissolves faster than troche, but slower than RDT.
- Do not swallow. Must dissolve in mouth to be effective, do not drink anything for at least 5 minutes after dissolving.



RAPID DISSOLVE TABLET (RDT)

- Round compressed tablet.
- Custom options available upon request.
- Fastest dissolving oral dosage form.
- Do not swallow, must dissolve in mouth to be effective, do not drink anything for at least 5 minutes after dissolving.



TROCHE

- Waxy lozenge.
- Custom options available upon request.
- Semi-solid dosage form, typically placed between the cheek and gum.
- Slowest dissolving oral dosage form.
- Requires refrigeration. (2-8 °C, 36-46 °F)



VAGINAL TROCHE

- Scored to provide ½ or ¼ dosing.
- Place troche just inside vaginal delivery device, insert in vagina and push end of device to insert medication.
- Requires refrigeration. (2-8 °C, 36-46 °F)



VAGINAL SUPPOSITORY

- Place suppository just inside vaginal delivery device, insert in vagina and push end of device to insert medication.
- Requires refrigeration. (2-8 °C, 36-46 °F)